Real-world evidence data reveals the physicians' perception on erenumab therapy in German headache centers



Scan to download a copy of thi presentation

Mirja Koch* 1, Katrin Schuh1, Charly Gaul2, Philipp Stude3, Andreas Straube4


¹Clinical Research Neuroscience, Novartis Pharma GmbH, Nuremberg, Germany ²Migraine and Headache Clinic, Koenigstein am Taunus, Germany; ³Neurologische Praxis Dr. Stude, Bochum, Germany, ⁴Ludwig-Maximillians-University, Munich, Germany

Migraine Trust Virtual Symposium, 3–9 October, 2020



- TELESCOPE was an online survey (July 2019 December 2019) of 45 headache centers in Germany:
 - Capturing the treating-physician's perspective on erenumab treatment with regards to therapy decision, patient profiles and quality of life (QoL) of the patients
 - Documentation of treatment effects and satisfaction with outcome of 10-20 individual episodic and chronic migraine patients with at least three months of erenumab treatment per site (in total 542 patients were documented)
- A concurrent project evaluated response and subjectiv judegement of patients (PERSICOPE; see MTIS DP-043)






• More than half of the physicians considered restricted QoL, monthly migraine days (MMD) and number of previous prophylactic treatments very important in deciding to initiate treatment with monoclonal antibodies



- Treating physicians observed reduced migraine burden and improved quality of life in more than 75% of their patients after treatment with erenumab
- Response rate and patient satisfaction were both reported as high

Figure 2. Erenumab effects and satisfaction










Figure 3. Erenumab effects on migraine specific parameters and onset of effect





- TELESCOPE provides us with real-world evidence data for erenumab in Germany
- Based on the assessment of treating physicians, erenumab reduces the burden of migraine and increases the quality of life in more than 75% of their patients, with an onset after first injection in a majority of the patients.